



**Nathan A. Mall, MD, CIME**  
The Orthopedic Center of St. Louis  
Cartilage Restoration Center of St. Louis  
14825 N. Outer 40, Suite 200  
Chesterfield, MO 63017  
P: 314.336.2555 F: 866.276.7014

Post-Operative Protocol-Lateral Epicondyle Debridement and Repair

Patient Name: \_\_\_\_\_ Date: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ Date of Surgery: \_\_\_\_\_

\_\_\_\_\_ Provide patient with home exercise program

**-Phase 1: Days 1-7**

- Position the extremity in a sling for comfort.
- Control edema and inflammation: Apply ice for 20 minutes two to three times a day.
- Gentle hand, wrist, and elbow ROM exercises. Exercises should be done in a pain-free ROM.
- Active shoulder ROM and Periscapular exercises
- Patient should minimize the frequency of any activities of daily living (ADLs) that stress the extensor tendon mechanism such as lifting, and combined joint movements (i.e. full elbow extension with wrist flexion). When lifting and/or performing activities with the surgical upper extremity it is advise to have the patient perform such tasks with their palm up to minimize work load extensor tendons. Consider pre-fab / custom wrist splint to minimize wrist extension activity if patient is acutely painful with such activities.
- Elbow pad for protection of incision site
- Education on work / activity modification.

**-Phase 2: Weeks 2-5**

- Discontinue sling.
- Begin passive range of motion (PROM). Passive motion should be continued and combined with active-assisted motion within end-range of patient's pain tolerance
- Gentle strengthening exercises with active motion and sub maximal isometrics.
- Edema and inflammation control: Continue to ice application 20 minutes two to three times a day. Tubigrip as needed.
- Scar management as needed.

**-Phase 3: Weeks 6-8**

- Advanced strengthening as tolerated to include weights or theraband. Focus should be on endurance training of wrist extensors (i.e. light weights, higher repetitions per set).
- ROM with continued emphasis on restoring full A/PROM.
- Edema and inflammation control with ice application for 20 minutes after activity.
- Modified activities in preparation for beginning functional training.
- Gentle massage along and against fiber orientation.
- Counterforce bracing to common extensor tendon of forearm. (Including education on proper use to avoid nerve compression.)

**-Phase 4: Weeks 8-12**

- Continue counterforce bracing for patient to completed ADLs and/or strengthening activity pain-free.
- Begin task-specific functional training.
- Return to higher-level work / recreational activities.

\_\_\_\_\_ Modalities as indicated

\_\_\_\_\_ days/week x \_\_\_\_\_ weeks

---

Nathan A. Mall, MD