



Nathan A. Mall, MD, CIME
The Orthopedic Center of St. Louis
Cartilage Restoration Center of St. Louis
14825 N. Outer 40, Suite 200; Chesterfield, MO 63017
P: 314.336.2555 F: 866.276.7014

Post-Operative Protocol- Shoulder Arthroscopy and Biceps Tenodesis

Patient Name: _____ Date: _____

Diagnosis: _____ Date of Surgery: _____

____ Provide patient with home exercise program

0-5 Weeks:

- Sling for first 4 weeks
- PROM → AAROM → AROM of elbow without resistance. This gives biceps tendon time to heal into new insertion site on humerus without being stressed
- Encourage pronation/supination without resistance
- Grip strengthening
- Maintain shoulder motion by progressing PROM → AROM without restrictions
- ROM goals: Full passive flexion and extension at elbow; full shoulder AROM
- No resisted motions until 4 weeks post-op
- Heat before PT sessions; other physical modalities per PT discretion

5-12 Weeks:

- D/C sling
- Begin AROM for elbow in all directions with passive stretching at end ranges to maintain or increase biceps/elbow flexibility and ROM
- At 6 weeks, begin light isometrics with arm at side for rotator cuff and deltoid; can advance to bands as tolerated
- At 6 weeks, begin scapular strengthening

3-12 Months:

- Only do strengthening 3x/week to avoid rotator cuff tendonitis
- Begin UE ergometer
- Begin eccentrically resisted motions, plyometric (ex. weighted ball toss), proprioception (ex. body blade), and closed chain exercises at 12 weeks.
- Begin sports related rehab at 3 months, including advanced conditioning
- Return to throwing and begin swimming at 3 months (see return to throwing program)
- Throw from pitcher's mound at 4 ½ months (see return to throwing program)
- Collision sports at 6 months

____ Other:

Modalities: _____ Electric Stimulation _____ Ultrasound _____ Heat before/Ice after

Frequency: _____ x/ week x _____ weeks

Nathan A. Mall, MD