



Nathan A. Mall, MD, CIME
The Orthopedic Center of St. Louis
Cartilage Restoration Center of St. Louis
14825 N. Outer 40, Suite 200; Chesterfield, MO 63017
P: 314.336.2555 F: 866.276.7014

Post-Operative Protocol- Posterior Labral Repair/Stabilization

Patient Name: _____ Date: _____

Diagnosis: _____ Date of Surgery: _____

_____ Provide patient with home exercise program

0-4 Weeks:

- Sling in neutral rotation for 4 weeks (padded abduction sling)
- Codman exercises, elbow and wrist ROM
- Wrist and grip strengthening

4-6 Weeks:

- Restrict to FF 90°/IR to stomach PROM → AAROM → AROM
- ER with arm at side as tolerated
- Begin isometrics with arm at side – FF/ER/IR/ABD/ADD
- Start scapular motion exercises (traps/rhomboids/lev. scap/etc.)
- No cross-arm adduction, follow ROM restrictions
- Heat before treatment, ice after treatment per therapist's discretion

6-12 Weeks:

- Increase ROM to within 20° of opposite side; no manipulations per therapist; encourage patients to work on ROM on a daily basis
- Once 140° active FF, advance strengthening as tolerated: isometrics → bands → light weights (1-5 lbs.); 8-12 reps/2-3 sets per rotator cuff, deltoid, and scapular stabilizers with low abduction angles
- Only do strengthening 3x/week to avoid rotator cuff tendonitis
- Closed chain exercises

3-12 Months:

- Advance to full ROM as tolerated
- Begin eccentrically resisted motions, plyometric (ex. Weighted ball toss), proprioception (es. body blade)
- Begin sports related rehab at 3 months, including advanced conditioning
- Return to throwing at 4 ½ months (see return to throwing program)
- Push-ups at 4 ½ - 6 months
- Throw from pitcher's mound at 6 months (see return to throwing program)

_____ Other:

Modalities: _____ Electric Stimulation _____ Ultrasound _____ Heat before/Ice after

Frequency: _____ x/ week x _____ weeks

Nathan A. Mall, MD